Extension of maximum likelihood estimation methods to mixed linear models

<u>Dário Ferreira</u>¹, Sandra Ferreira¹, Célia Nunes¹, and João T. Mexia²

¹University of Beira Interior, Covilhã, Portugal ²New University of Lisbon, Portugal

Abstract

We extend and compare Maximum Likelihood Estimation methods, that usually are used for estimating the parameters in unbalanced random linear models, to unbalanced mixed linear models.

Keywords

Linear mixed models, Variance components, Maximum likelihood estimation methods.

References

Ferreira, D., Ferreira, S., Nunes, C., and Mexia, J.T. (2009). Adjustment of normal mixed models through triple minimization. *InterStat.*

Fonseca, M., Carvalho, M., Oliveira, M., and Mexia, J.T. (2008). A reduction technique for conducting inference in mixed models. *LinStat'2008* - the talk.

Fonseca, M. and Mexia J.T. (2007). Inference in mixed models with stochastic search. *Proceedings of the 56th Session of the International Statistics Institute 2007.*

Hartley, H.O. and Rao, J.N.K. (1967). Maximum likelihood estimation for the mixed analysis of variance model. *Biometrika* 54, 93–108.

Patterson, D. and Thompson, R. (1974). Maximum likelihood estimation of components of variance. *Proceedings of the 8th International Biometric Conference*, 197–207.